arXiv:hep-th/0111108v1 12 Nov 2001

Quantum Properties of Solitons in $U(\phi) = \phi^2 \ln^2(\phi^2)$ and $U(\phi) = \phi^2 \cos^2(\phi^2)$ Models

Gabriel H. Flores[1] *,and N.F. Svaiter.[2] †

[1] *Centro Brasileiro de Pesquisas Fisicas-CBPF,*
Rua Dr.Xavier Sigaud 150, 22290-180 Rio de Janeiro, RJ, Brazil
[2] *Center of Theoretical Physics,*
Massachusetts Institute of Technology,
Cambridge, MA 02139, USA.

February 1, 2008

Abstract

Recently we constructed two new $(1+1)$-dimensional scalar field theory models that posses solitonic solutions. They are the $U(\phi) = \phi^2 \ln^2(\phi^2)$ and the $U(\phi) = \phi^2 \cos^2(\phi^2)$ models . The first quantum corrections for these models are given by exactly solvable Schrodinger equations. In this paper we first examine the quantum meaning of the solitonic solutions and study the scattering of the mesons by the quantum soliton at order $\hbar$. Finally we give a finite expression for the soliton masses of both models and evaluate such expression approximately in the case of the second model.

PACS number(s):11.30.Pb, 03.65.Fd, 11.10.Ef.

---

[*] E-mail: gflores@lafex.cbpf.br

[†] Permanent address: Centro Brasileiro de Pesquisas Fisicas, Rio De Janeiro, RJ 22290-180, Brazil. E-mail: svaiter@lns.mit.edu, nfuxsvai@cbpf.br

1 Introduction

In a recent work [1] the authors have constructed two new $(1+1)$-dimensional scalar field theory models starting from analyticaly solvable Schrodinger equations. Starting from the Morse potential we have obtained the model with density potential given by

$$U(\phi) = \frac{m^2}{8}\phi^2 \ln^2\left(\frac{\alpha^2\phi^2}{9m^4}\right) , \tag{1}$$

and starting from the Scarf II hyperbolic potential we obtained the model with density potential given by

$$U(\phi) = \frac{1}{2}m^2 B^2 \phi^2 \cos^2\left[\frac{1}{2B}\ln\left(\frac{\alpha^2\phi^2}{9m^4}\right)\right] . \tag{2}$$

(In ref.[1] we fixed B=1/2) The profiles of these density potentials given by eqs. (1) and (2) are showed in fig.(1) and fig.(2) respectively. In that follows we refer to the models with density potentials given by equations (1) and (2) as models I and II respectively.

For the model I it is easy to obtain the kink and anti-kink like solutions. They are given by

$$\phi_c(x) = \pm\phi_0 \exp(-e^{\pm m(x-x_0)}) , \tag{3}$$

where $\phi_0 = 3m^2/\alpha$ [1]. In this case we have two kinks and two anti-kinks respectively that lie between two of the three degenerate trivial vacua that are showed in fig.(1). The energy of these classical configurations are all the same and are given by

$$\begin{aligned} H[\phi_c] &= \int_{-\infty}^{\infty} dx \left(\frac{1}{2}\left(\frac{d\phi_c}{dx}\right)^2 + \frac{m^2}{8}\phi_c^2\left[\ln\left(\frac{\alpha^2\phi_c^2}{9m^4}\right)\right]^2\right) \\ &= \frac{9m^6}{2\alpha^2}\int_{-\infty}^{\infty} dx e^{\pm mx}\exp(-2e^{\pm mx}) \\ &= \frac{9m^5}{4\alpha^2} . \end{aligned} \tag{4}$$

We can make a Taylor expansion around one of the trivial vacua and we obtain

$$U(\varphi) = \frac{1}{2}m^2\varphi^2 \pm \frac{\alpha}{6}\varphi^3 - \frac{1}{216}\varphi^4 + \frac{1}{14580}\frac{\alpha^4}{m^6}\varphi^6 + ... \tag{5}$$

where we have shifted the field as $\phi = (\varphi \pm \phi_0)$. We remark that an expansion around the point $\phi = 0$ is not possible because with the exception of the first derivative all the derivatives of $U(\phi)$ are infinite at this point. This means that it is not possible to have perturbative dynamics around the trivial vacuum $\phi = 0$. Note that this model possesses a discrete symmetry $U(\phi) = U(-\phi)$ that is broken by the perturbative vacua $\pm\phi_0$.

Model II has infinitely degenerate trivial vacua at the points $\phi = \pm\phi_n$ with ϕ_n given by

$$\phi_n = \frac{3m^2}{\alpha} \exp\left(\frac{2n+1}{2}\pi B\right) \ , \quad n = 0, \pm 1, \pm 2, .. \tag{6}$$

The kinks and anti-kinks that lie between two of these trivial vacua are given by

$$\phi_c(x) = \pm\frac{3m^2}{\alpha} \exp\left(n\pi B \pm B \tan^{-1}(\sinh(mx))\right) \ , \quad n = 0, \pm 1, \pm 2, .. \tag{7}$$

where the solutions with $(\pm)$ signs in the exponents correspond to the kinks and anti-kinks solutions respectively for each value of n and for each sign that appears in front. Also this model possesses the discrete symetry $U(-\phi) = U(\phi)$ that is broken by the trivial vacua. Notice that $\lim_{n\to\infty} \phi_n = 0$, *i.e*, the density potential oscillates infinite times between some finite value of ϕ and zero. Also observe that the distance between the trivial vacua grows out exponentially. We can also obtain the energy for these configurations, they are given by

$$\begin{aligned} H[\phi_c] &= \int_{-\infty}^{\infty} \left(\frac{1}{2}\left(\frac{d\phi_c}{dx}\right)^2 + \frac{m^2}{8}\cos^2\left[\ln\left(\frac{\alpha^2\phi^2}{9m^4}\right)\right]\right) \\ &= \frac{9m^6 B^2 e^{2Bn\pi}}{\alpha^2} \int_{-\infty}^{\infty} dx \frac{\exp(\pm 2B \arctan(\sinh(mx)))}{\cosh^2(mx)} \\ &= \frac{9m^5 B^2 e^{2Bn\pi}}{\alpha^2} I(B) \end{aligned} \tag{8}$$

where $I(B)$ is given by

$$I(B) = \int_{-1}^{1} ds e^{2B \arcsin(s)} \tag{9}$$

and can be evaluated numericaly for each value of B. In eq. (8) we can see that the energy associated with these static configurations are all different. As in the case of the model I, we can make a Taylor expansion around one of the trivial vacua and we obtain

$$\begin{aligned} U(\varphi) &= \frac{m^2}{2}\varphi^2 \pm \frac{\alpha}{6} e^{-\frac{1}{2}(2n+1)\pi B}\varphi^3 - \frac{(1+4/B^2)}{9(4!)}\frac{\alpha^2}{m^2} e^{-(2n+1)\pi B}\varphi^4 \\ &\qquad + \frac{(4+20/B^2+16/B^4)}{(81)6!}\frac{\alpha^4}{m^6} e^{-2(2n+1)\pi B}\varphi^6 + ... \end{aligned} \tag{10}$$

Here we have redefined $\phi = (\varphi \pm \phi_n)$. From the Taylor expansion we can see a peculiarity of this model. When $n \to \infty$ all the terms vanish except the quadratic term, *i.e*, the perturbative sector around the trivial vacuum for $n \to \infty$ becomes a free theory. Also as in the model I we have to stress that when $n \to -\infty$ the Taylor expansion given by eq.(10) is meaningless, *i.e*, there is no perturbative dynamics around the trivial vacuum $\phi = 0$.

In the next section following Jackiw[?] we will use the Kerman-Klein method to analyse the quantum properties of the solitonic solutions.

2 Quantum meaning of the solitonic solutions: the Kerman-Klein method

To analyse the quantum properties of the solitonic solutions, we use the Kerman-Klein method [2], [3]. For such purpose we postulate that in addition to the perturbative states (that we call meson states), there are other particle states, the quantum soliton states (in the literature they are called the baryons states). The one-soliton state $|E>$ are energy and momentum eigenstates

$$\begin{aligned} \hat{H}|E> &= E|E> \ , \\ \hat{P}|E> &= P|E> \ , \end{aligned} \tag{11}$$

where $E = \sqrt{P^2 + M^2}$ and M is the mass of the quantum soliton state. Further it is postulated that for small coupling constant (α small) the quantum soliton is very heavy, $i.e$, M is taken to be

$$M \approx \mathcal{O}(\alpha^{-2}) \ . \tag{12}$$

In addition we postulate that there are one-soliton multi-meson states, $|P, K_1, K_2 ... >$ where P is the total momentum and k_i is the asymptotic meson momentum. Also we postulate the stability of the quantum solitonic state, $i.e$, that matrix elements as

$$< \text{meson state, soliton state}|\hat{\phi}\hat{\phi}...|\text{meson state, no soliton state} >= 0 \tag{13}$$

vanish. In model I we can see that there are two types of soliton states, $i.e$, the corresponding to the ($\pm$) signs that appear in the exponential of eq.(3). These states are called soliton and anti-solitons states respectively. For the model II we have an infinite number of soliton and anti-soliton states, labeled by the index n of eq. (7). In both cases we are considering that the soliton and anti-soliton states corresponding to the ($\pm$) signs that appear in front to the eqs. (3) and (7) are the same. This is justified by the discrete symetry $U(\phi) = U(-\phi)$ in both cases. The sectors of the Hilbert state constructed taking into acount the soliton and anti-soliton states are called the soliton and anti-soliton sectors respectively. Finally we postulate that the soliton sectors do not comunicate with the anti-solitonic sectors. Now we can verify using the above postulates and self-consistence that

$$< P'; K_1', ..., K_n'|\hat{\phi}|P; K_1, ..., K_m' >_c \ \approx \mathcal{O}(\alpha^{n+m-1}) \ , \tag{14}$$

where the subscript c denotes connected part and $\hat{\phi}$ is a Heisenberg field operator. For this purpose we use the equation of motion for the field operator ϕ,

$$\frac{\partial^2 \hat{\phi}}{\partial t^2} - \frac{\partial^2 \hat{\phi}}{\partial x^2} = -U'(\hat{\phi}) \ . \tag{15}$$

In first place we will compute $f(P', P) =< P'|\hat{\phi}|P >$ to the order $\mathcal{O}(\alpha^{-1})$. In both models we can make a taylor expansion for $U(\hat{\phi})$ around one of the trivial vacua. It is easy to show that such expression in both cases have the following form

$$U(\hat{\phi}) = \sum_{l=2}^{\infty} a_l \alpha^{l-2} (\hat{\phi} - \phi_0)^l \ . \tag{16}$$

In eq.(16) ϕ_0 denotes one of the trivial vacua and the coefficient a_l does not depend on α. Using eq.(16) in eq.(15) we have

$$\frac{\partial^2 \hat{\phi}}{\partial t^2} - \frac{\partial^2 \hat{\phi}}{\partial x^2} = -\sum_{l=2}^{\infty} l a_l \alpha^{l-2} (\hat{\phi} - \phi_0)^{l-1} \ . \tag{17}$$

Using the Heisenberg equations of motion

$$\begin{aligned} \frac{\partial \hat{\phi}}{\partial t} &= i[\hat{H}, \hat{\phi}] \\ \frac{\partial \hat{\phi}}{\partial x} &= -i[\hat{P}, \hat{\phi}] \end{aligned} \tag{18}$$

twice in eq. (17) we obtain

$$\Big((P - P')^2 - (E - E')^2\Big) f(P', P) = -\sum_{l=2}^{\infty} l a_l \alpha^{l-2} < P'|(\hat{\phi} - \phi_0)^{l-1}|P > \ . \tag{19}$$

Each element $f^{l-1}(P', P) =< P'|(\hat{\phi} - \phi_0)^{l-1}|P >$ can be written as

$$\begin{aligned} f^{l-1}(P', P) \ &= \ \sum_{m_i} < P'|(\hat{\phi} - \phi_0)|m_1 >< m_1|(\hat{\phi} - \phi_0)|m_2 >< m_2|... \\ &\qquad ...|m_{l-2} >< m_{l-2}|(\hat{\phi} - \phi_0)|P > \ , \end{aligned} \tag{20}$$

where the sum is over the complete basis $|m_i >$, *i.e*, over the soliton states, the soliton one meson states, the soliton two meson states , etc. In eq.(19) the left hand side is of the order $\mathcal{O}(\alpha^{-1})$. For consistency we have to take the right hand side also of this order. In order to make this possible we have to take each $f^{l-1}(P',P)$ term of the order $\mathcal{O}(\alpha^{-(l-1)})$. Then in eq.(20) we have to retain only the sum over the one solitonic (or one anti-solitonic, if it were the case) states. Then we have for $f^{l-1}(P',P)$:

$$\int \frac{dP_1}{2\pi}\frac{dP_2}{2\pi}...\frac{dP_{l-2}}{2\pi}[f(P',P_1)-2\pi\phi_0\delta(P'-P_1)][f(P_1,P_2)-2\pi\phi_0\delta(P_1-P_2)]... \\ ...[f(P_{l-2},P)-2\pi\phi_0\delta(P_{l-2}-P)]\ . \tag{21}$$

Replacing eq.(21) in eq.(19) we obtain an integral equation for $f(P',P)$,

$$\left((P-P')^2-(E-E')^2\right)f(P',P) = -\sum_{l=2}^{\infty} l a_l \alpha^{l-2}\int\frac{dP_1}{2\pi}...\frac{dP_{l-2}}{2\pi} \\ [f(P',P_1)-2\pi\phi_0\delta(P'-P_1)]...[f(P_{l-2},P)-2\pi\phi_0\delta(P_{l-2}-P)]\ . \tag{22}$$

To solve eq.(22) we note that at this order $(E-E')\approx 0$. This can be seen using $E=\sqrt{P^2+M^2}\approx M+\frac{P^2}{2M}$ and noting that the second term is of the order $\mathcal{O}(\alpha^2)$, *i.e*, at this order the energy is independent of the momentum. This is compatible with the assumption that the quantum soliton is very heavy. Also using the Lorentz invariance of $f(P',P)$ we can set at this order $f(P',P)=f(P-P)$. Using these facts eq. (22) can be reduced to

$$\begin{aligned}(P-P')^2 f(P',P) &= -\sum_{l=2}^{\infty} l a_l\alpha^{l-2}\int\frac{dP_1}{2\pi}...\frac{dP_{l-2}}{2\pi}[f(P'-P_1)-2\pi\phi_0\delta(P'-P_1)]... \\ &\quad ...[f(P_{l-2}-P)-2\pi\phi_0\delta(P_{l-2}-P)]\ .\end{aligned} \tag{23}$$

We can solve eq.(23) in terms of a Fourier transformed function $\phi(x)$

$$f(P'-P)=\int dx\,\exp[i(P'-P)x]\phi(x)\ . \tag{24}$$

In term of $\phi(x)$ eq. (23) is

$$\begin{aligned}\phi''(x) &= \sum_{l=2}^{\infty} l a_l\alpha^{l-2}\left(\phi(x)-\phi_0\right)^{l-1} \\ &= U'(\phi).\end{aligned} \tag{25}$$

We recognize eq.(25) as the classical equation of motion for ϕ_c. Therefore, at the order $\mathcal{O}(\alpha^{-1})$, we obtain for the matrix elements for the quantum field operator between one soliton (or one anti-soliton) states:

$$< P'|\hat{\phi}|P >=\int dx\,\exp[i(P'-P)x]\phi_c(x)\ , \tag{26}$$

where $\phi_c(x)$ is given by equation (3) or (7). Since in both models I and II $\phi_c(x)$ is proportional to $1/\alpha$ one verifies in eq.(26) that $< P'|\hat{\phi}|P >$ is $\mathcal{O}(\alpha^{-1})$. We can compute at first leading order the mass of the quantum solitonic state using that at leading order $\hat{H}|P >= \sqrt{P^2+M^2}|P >\approx M|P >$. The energy operator is

$$\hat{H} = \int dx \left(\frac{1}{2}\partial_\mu\hat{\phi}\partial_\mu\hat{\phi} + U(\hat{\phi})\right) \ . \tag{27}$$

Replacing eq.(16) in eq. (27) we obtain

$$\begin{aligned} < P|\hat{H}|P > &= \int dx \left(\frac{1}{2} < P|\partial_\mu\hat{\phi}\partial_\mu\hat{\phi}|P > + \sum_{l=2}^{\infty} a_l \alpha^{l-2} < P|(\hat{\phi}-\phi_0)^l|P >\right) \\ &= \int dx \left(\frac{1}{2}\sum_{m_1} < P|\partial_\mu\hat{\phi}|m_1 >< m_1|\partial_\mu\hat{\phi}|P > + \sum_{l=2}^{\infty}\sum_{m_i} a_l\alpha^{l-2}\right. \\ &\quad \left. < P|(\hat{\phi}-\phi_0)|m_1 >< m_1|...|m_{l-1} >< m_{l-1}|(\hat{\phi}-\phi_0)|P >\right) \end{aligned} \tag{28}$$

and since the left hand side of eq.(28) is of $\mathcal{O}(\alpha^{-2})$ order we can see that in the sum over m_i we have only to retain the sum over the one soliton states and using eq.(18) in the first term of the right hand side of eq.(28) we obtain

$$\begin{aligned} < P|\hat{H}|P > &\approx 2\pi\delta(P-P)M \\ &= 2\pi\delta(0)\int dx \left(\frac{1}{2}\left(\frac{d\phi_c}{dx}\right)^2 + U(\phi_c)\right) \ . \end{aligned} \tag{29}$$

Then we have for the soliton mass at leading order

$$M = \int dx \left(\frac{1}{2}\left(\frac{d\phi_c}{dx}\right)^2 + U(\phi_c)\right) \ , \tag{30}$$

that is, it is equal to the energy of the classical static configuration. For model I it is given by the eq.(4) and for model II it is given by eq.(8). In both cases M is $\mathcal{O}(\alpha^{-2})$ confirming eq.(12). Note that in model I the two different soliton states have the same mass at this order. On the other hand the mass of the soliton states of the model II are different. Note also that the soliton mass becomes infinite when $n \to \infty$. In order to compute $f(P',P)$ to the next order we have to retain in eq.(20) the sum over soliton one meson states $|P,k >$, then we will obtain the next term for $f^{l-1}(P',P)$:

$$\begin{aligned} &\int \frac{dP_1}{2\pi}\frac{dk_1}{2\pi}\frac{dP_2}{2\pi}...\frac{dP_i}{2\pi}...\frac{dP_{l-2}}{2\pi} < P'|(\hat{\phi}-\phi_0)|P_1;k_1 >< P_1;k_1|(\hat{\phi}-\phi_0)|P_2 >< P_2|... \\ &\qquad |P_i >< P_i|(\hat{\phi}-\phi_0)|P_{i+1} >< P_{i+1}|...|P_{l-2} >< P_{l-2}|(\hat{\phi}-\phi_0)|P > +... \\ &... + \int \frac{dP_1}{2\pi}\frac{dP_2}{2\pi}...\frac{dP_i}{2\pi}\frac{dk_i}{2\pi}...\frac{dP_{l-2}}{2\pi} < P'|(\hat{\phi}-\phi_0)|P_1 >< P_1|(\hat{\phi}-\phi_0)|P_2 >< P_2|... \end{aligned}$$

$$
\begin{aligned}
&|P_i;k_i>< P_i;k_i|(\hat{\phi}-\phi_0)|P_{i+1}>< P_{i+1}|...|P_{l-2}>< P_{l-2}|(\hat{\phi}-\phi_0)|P> +...\\
&...+\int\frac{dP_1}{2\pi}\frac{dP_2}{2\pi}\frac{dP_i}{2\pi}...\frac{dP_{l-2}}{2\pi}\frac{dk_{l-2}}{2\pi}< P'|(\hat{\phi}-\phi_0)|P_1>< P_1|(\hat{\phi}-\phi_0)|P_2>< P_2|...\\
&|P_i>< P_i|(\hat{\phi}-\phi_0)|P_{i+1}>< P_{i+1}|...|P_{l-2};k_{l-2}>< P_{l-2};k_{l-2}|(\hat{\phi}-\phi_0)|P> \ . \qquad (31)
\end{aligned}
$$

Since $f_k(P_1,P_2) =< P_1|\hat{\phi}|P_2,k >$ is of the order $O(\alpha^0)$ then the contribution for $f^{l-1}(P',P)$ given by eq.(31) is of the order $O(\alpha^{3-l})$ and from eq.(19) we can see that the expression (31) is the $O(\alpha^1)$ contribution for $f(P',P)$. We can see that to compute $f(P',P)$ to the next order we need to know $f_k(P_1,P_2)$ at lower order. Also if one computes the energy, to the next order we will need $f_k(P_1,P_2)$. Then our next task is to evaluate $f_k(P_1,P_2)$. Using eq.(17) and eq.(18) we obtain

$$
\left((P_2-P_1)^2-(E_2+\omega(k)-E_1)^2\right)f_k(P_1,P_2)=-\sum_{l=2}^{\infty}la_l\alpha^{l-2}< P_1|(\hat{\phi}-\phi_0)^{l-1}|P_2;k> \ . \qquad (32)
$$

We can write $f_k^{l-1}(P_1,P_2) =< P_1|(\hat{\phi}-\phi_0)^{l-1}|P_2;k>$ as in eq.(20)

$$
\begin{aligned}
f_k^{l-1}(P_1,P_2) &= \sum_{m_i}< P_1|(\hat{\phi}-\phi_0)|m_1>< m_1|(\hat{\phi}-\phi_0)|m_2>< m_2|...\\
&\qquad ...|m_{l-2}>< m_{l-2}|(\hat{\phi}-\phi_0)|P_2;k> \ . \qquad (33)
\end{aligned}
$$

The left hand side of eq.(32) is of the order $O(\alpha^0)$ then $f_k^{l-1}(P_1,P_2)$ need to be of the order $O(\alpha^{2-l})$. If we retain in eq.(33) the sum only over soliton states it can be seen that $f_k^{l-1}(P_1,P_2)$ is of required order $O(\alpha^{2-l})$. But if we keep in the sum over m_{l-2} soliton one meson states (in the other m_i's we sum over soliton states only) using

$$
< P_{l-2};k_{l-2}|\hat{\phi}|P_2;k>=2\pi\delta(k_{l-2}-k)< P_{l-2}|\hat{\phi}|P_2>+< P_{l-2};k_{l-2}|\hat{\phi}|P_2;k>_c \qquad (34)
$$

and retaining only the disconected term we can see that also such term will be of the required $O(\alpha^{2-l})$. Nextly we can retain in the sums over m_{l-2} and m_{l-3} one meson soliton states and retaining as above only the disconected terms it can be seen that also the term obtained will be of the order $O(\alpha^{2-l})$. Next we retain in the sum over m_{l-2}, m_{l-3} and m_{l-4} soliton one meson states, next in m_{l-2}, m_{l-3} m_{l-4} and m_{l-5} and so on. All these terms will be of the order $O(\alpha^{2-l})$. We see that there is one term with origin in the sum over soliton states, and $(l-2)$ terms from summing over soliton one meson states, that is we have $(l-1)$ terms of the order $O(\alpha^{2-l})$. At this order still $E=M$. And using a Fourier transformed function $\phi_k(x)$

$$
f_k(P_2,P_1)=\int dx\exp[i(P_2-P_1)x]\phi_k(x) \qquad (35)
$$

we can prove that all the $(l-1)$ terms give the same contribution in eq.(32). Then, we obtain the following equation for $\phi_k(x)$

$$\begin{aligned}\left[-\frac{d^2}{dx^2}-\omega^2(k)\right]\phi_k(x) &= -\sum_{l=2}^{\infty} l(l-1)a_l\alpha^{l-2}(\phi_c(x)-\phi_0)^{l-2}\phi_k(x) \\ &= -U''(\phi_c(x))\phi_k(x)\ , \end{aligned} \tag{36}$$

that is, we obtain a Schrodinger like equation for $\phi_k(x)$

$$\left[-\frac{d^2}{dx^2}+U''(\phi_c(x))\right]\phi_k(x)=\omega^2(k)\phi_k(x). \tag{37}$$

Of course eq.(37) has discrete and continuum eigenvalues $\omega^2(k)$. The discrete eigenvalues can be interpreted as excited states of the soliton states. The continuum eigenvalues are $\omega(k)=\sqrt{k^2+m^2}$ (where m is the mass of the mesons) and the eigevalues asymptoticaly behave like e^{ikx}, $i.e$, meson plane waves. That is, at this order the scattering of mesons by solitons are described effectively by the scattering problem given by eq.(37). It is easy to show that $d\phi_c/dx$ is the eigenfuntion of (37) with eigenvalue zero. It was shown [**?**] that these eigenfuntions can be disregarded from the complete basis $\phi_k(x)$ consistently with Poincare invariance and the conmmutation relation between the field operator and its canonically conjugate momentum. The zero mode solution of eq.(37) has no physical meaning, it gives only the first corrections to the movement of the soliton. To show the above one needs to write $\phi_k(x)$ in terms of normalized eigenfuntions $\psi_k(x)$, as $\phi_k(x)=\psi_k(x)/\sqrt{2\omega(k)}$ and then we have for $f_k(P_2,P_1)$

$$f_k(P_2,P_1)=\int dx\exp[i(P_2-P_1)x]\frac{\psi_k(x)}{\sqrt{2\omega(k)}} \tag{38}$$

where now we understand that the zero modes are excluded. Using eq.(38) in expression (31) one can prove that $f(P',P)$ is given by eq.(24) but now with $\phi(x)$ given by

$$\phi''(x)=U'(\phi)+\frac{1}{2}G(x,x)U'''(\phi)\ , \tag{39}$$

where

$$G(x,y)=\sum{}'\frac{\psi_k^\star(x)\psi_k(y)}{2\omega(k)}\ , \tag{40}$$

where the prime in the sum indicate that the zero mode is excluded. Also we can compute the mass of the soliton to the next order retaining in (28) the contribution of soliton one meson states, obtaining

$$M=H[\phi_c]+\frac{1}{2}\sum_k\omega(k)\ . \tag{41}$$

3 The scattering of mesons by solitons

As stated above to the scattering of mesons by solitons we have to solve the one dimensional scattering problem for the Schrodinger equation given by eq.(37). For the model I the eq.(37) is given by

$$\left[-\frac{d^2}{dx^2}+m^2(e^{\pm 2mx}-3e^{\pm mx}+1)\right]\phi_k(x)=\omega^2(k)\phi_k(x)\ , \tag{42}$$

where the signs ± are respectively for the soliton anti-soliton sectors. Here we solve explicitly the + case. In fig.(3) we show this potential for the + sign. Solving eq.(42) one finds that there is only one discrete eigenfuntion [4]

$$\psi_0(x)=2\sqrt{m}e^{mx}\exp(-e^{mx})\ , \tag{43}$$

with zero eigenvalue $\omega_0^2=0$, and continuous eigenfuntions with eigenvalues going from m^2 to infinity. As we stated in preceding section the mode zero has no physical interpretation. Because there is no more discrete eigenfuntions, in this case the soliton state can not exist in a excited state. The continuum eigenfuntions are interpreted as the scattering of mesons by the soliton. One can see from the potential profile showed in fig.(3) that mesons coming from $x=-\infty$ can not reach $x=\infty$, they are absolutely reflected by the soliton going back again to $x=-\infty$. Then one can espect to find for the reflection and transmitions amplitudes $R=e^{i\delta(k)}$ and $T=0$ respectively, with $\delta(k)$ a real function. In order to find δ we have to solve eq.(42) with the following boundary conditions

$$\psi_k(x\to-\infty)=Ae^{ikx}+Be^{-ikx}\ , \tag{44}$$

and

$$\psi_k(x\to\infty)=0\ . \tag{45}$$

To compute R we have to compute the relation between the values A and B. For this end we make the following change of variables

$$\xi=2e^{mx},\quad \psi_k(x)=e^{-\xi/2}\xi^{ik/m}\chi(\xi)\ , \tag{46}$$

in eq.(42) and we obtain the following equation for $\chi(\xi)$

$$\xi\chi''(\xi)+(2ik/m+1-\xi)\chi'(\xi)+(1-ik/m)\chi(\xi)=0\ , \tag{47}$$

a confluent hypergeometric equation, whose solution is given in terms of hypergeometric functions[4]

$$\chi(\xi)=c_1F(ik/m-1,2ik/m+1,\xi)+c_2\xi^{-2ik/m}F(-ik/m-1,1-2ik/m,\xi)\ . \tag{48}$$

Using eq.(48) in eq.(46) we obtain for $\psi_k(x)$

$$\psi_k(x) = e^{-\xi/2}\left[c_1\xi^{ik/m}F(ik/m-1,2ik/m+1,\xi)+c_2\xi^{-ik/m}F(-ik/m-1,1-2ik/m,\xi)\right] \; , \tag{49}$$

where we can see that for $\xi \to 0$ ($x \to -\infty$) this solution behaves like eq.(44). Using the asymptotic value for $F(\alpha,\gamma,\xi)$ [5]

$$F(\alpha,\gamma,\xi\to\infty) = \frac{\Gamma(\gamma)}{\Gamma(\alpha)}e^{\xi}\xi^{\alpha-\gamma} \; , \tag{50}$$

it is easy to see that solution given by eq.(49) behaves as eq.(45) only if

$$c_1 = \frac{\Gamma(ik/m-1)\Gamma(-2ik/m)}{\Gamma(-1-ik/m)\Gamma(2ik/m)}c_2 \; . \tag{51}$$

Now using eq.(51) in eq.(49) and taking the limit $x \to -\infty$ we obtain

$$\psi_k(x\to-\infty) = c_2\left[\frac{\Gamma(ik/m-1)\Gamma(-2ik/m)}{\Gamma(-1-ik/m)\Gamma(2ik/m)}2^{ik/m}e^{ikx}+2^{-ik/m}e^{-ikx}\right] \tag{52}$$

from wich we obtain for the reflection amplitude:

$$R(k) = B/A = 2^{-2ik/m}\frac{\Gamma(-1-ik/m)\Gamma(2ik/m)}{\Gamma(-1+ik/m)\Gamma(-2ik/m)} \; . \tag{53}$$

And $\delta(k)$ is obtained as the argument of the above expression,

$$\delta(k) = Arg\left(2^{-2ik/m}\frac{\Gamma(-1-ik/m)\Gamma(2ik/m)}{\Gamma(-1+ik/m)\Gamma(-2ik/m)}\right) \; . \tag{54}$$

Note that as expected $|R(k)| = 1$. The case ($-$) in eq.(42) follows identicaly as the above case. The mesons caming from $x \to \infty$ never reach $x \to -\infty$, they are totally reflected by the soliton back again to $x \to \infty$. The reflection amplitude is given as in the ($+$) case by eq.(53).

For model II the Schrodinger equation (37) is given by

$$\left[-\frac{d^2}{dx^2}+m^2\left(1+\frac{(B^2-2)}{\cosh^2(mx)}\mp 3B\frac{\tanh(mx)}{\cosh(mx)}\right)\right]\phi_k(x) = \omega(k)^2\phi_k(x) \; , \tag{55}$$

where the signs ($\mp$) are related to the soliton and antisoliton sectors respectively. We show this potential in fig.(4) for the $-$ case.This equation have one discrete eigenvalue[6]

$$\psi_0(x) = N\frac{\exp(\pm 2B\tan^{-1}(\sinh(mx)))}{\cosh(mx)} \; , \tag{56}$$

where N is a normalization factor, and the continuum eigenfuntions with eigenvalues given by $\omega^2(k) = k^2+m^2$. As in the model I there are not excited soliton states. From fig.(4) one can expect the following behaviour of the scattering of mesons by the soliton (anti-soliton): Mesons with momenta k coming from $x = -\infty$ are scattered by the soliton (anti-soliton) in mesons with momenta k that reach $x = \infty$ (transmited mesons) and in mesons with momenta k that go back again to $x = -\infty$ (reflected mesons). To find the reflection and transmission amplitudes we have to solve eq.(55) with the adequate boundary conditions. The solutions of eq.(55) can be written in terms of hypergeometric functions and using the asymptotic properties of these hypergeometric functions one can find easily the following for the reflection and transmision amplitudes[7]

$$R(k) = \pm T(k) \frac{\sinh(\pi B)}{\cosh(\pi k/m)} \ , \tag{57}$$

and

$$T(k) = \frac{\Gamma(-1 - ik/m)\Gamma(2 - ik/m)\Gamma(\frac{1}{2} \mp iB - ik/m)\Gamma(\frac{1}{2} \pm iB - ik/m)}{\Gamma(-ik/m)\Gamma(1 - ik/m)\Gamma^2(\frac{1}{2} - ik/m)} \ . \tag{58}$$

Note that $T(k)$ is the same for the scattering of mesons by the soliton and anti-soliton states. $R(k)$ only differ by a phase equal to 2π, *i.e*, the scattering of mesons coming from $x = -\infty$ by the soliton state is the same as the scattering of mesons coming from $x = \infty$ by the anti-soliton state. One can also verify that $(|R(k)|^2 + |T(k)|^2) = 1$.

4 Quantum corrections for the soliton mass

The first quantum correction for the soliton mass is given by the sum in eq.(41) over all the eigenvalues of the Schrodinger equation. Such expression is divergent and if we subtract the zero point energy associated with one of the trivial vacua the divergence is rendered logarithmic. Then we need to renormalize such expression in order to obtain a finite expression. But since we are in $1 + 1$ it is well known that all divergences can be eliminated by normal ordering the field operator. Using this fact Cahill *et al* in reference [8] (see also [9]) found an expression free of divergences for the first quantum correction for the soliton mass. Expressed in terms of the normalized eigenfuntions of the Schrodinger equation this finite expression is given by[9]

$$\delta M = -\frac{1}{4} \sum_n \int_{-\infty}^{\infty} dq (\omega_n^2 \omega_q^{-1} - 2\omega_n + \omega_q) |\tilde{\psi}_n(q)|^2 \ , \tag{59}$$

where

$$\tilde{\psi}_n(q) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dx \exp(iqx) \psi_n(x) \ . \tag{60}$$

and $\omega_q = \sqrt{q^2 + m^2}$. Although in both models I and II we know explicitly all the eigenvalues and eigenfuntions it is still very hard to evaluate the expression given by eq.(59). But for model II there is a situation in which one can obtain δM aproximatelly. This situation is when B is very small. In this case from eq.(57) we see that $R \approx 0$, *i.e*, the Schrodinger equation associated to the model II is approximatelly reflectionless. And as was shown in ref.[8] for reflectionless potentials, δM given by eq.(59) can be expressed in terms of the discrete eigenvalues only, *i.e*,

$$\delta M = -\frac{m}{\pi} \sum_i (\sin\theta_i - \theta_i \cos\theta_i) \ , \tag{61}$$

where $\theta_i = \arccos(\omega_i/m)$. Since For model II there is only one (zero) discrete eigenvalue we obtain $\delta M = -\pi/m$. Then taking in equation (8) $I(B) \approx 2$ we can write for B small the following expression for the soliton masses asociated to model II

$$M = \frac{18 m^5 B^2 e^{2Bn\pi}}{\alpha^2} - \frac{m}{\pi} \ . \tag{62}$$

Since the above expression is valid for B small one would naively discard the first term. This can not be made since α is a small parameter.

5 Conclusions

In this paper we have analyzed the quantum properties of solitons in (1+1) field theory models with density potential given by equations (1) and (2) that we called models I and II respectively. Also we analyzed the scattering of mesons by solitons in both models at $\hbar$ order. For model II we computed aproximatelly the first quantum corrections for the mass of the soliton quantum state. It is interesting to remark that in model II the perturbative sector around the vacuum for large n is a free theory (see eq.(10)) but the solitonic sector is not. The mesons are scattered by the quantum soliton state. Then the lesson that we have learned is that if a theory is free in one sector there can be other sector(s) where it is not.

Acknowledgements

GHF and NFS are partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq (Brazil).

References

[1] G. Flores and N.F. Svaiter, Pre-print CBPF-NF-014-01, hep-th/0107043, submitted to Int.J.Mod.Phys.**A**.

[2] R. Jackiw and C. Rebbi, Phys.Rev. **D13**, 3398 (1976).

[3] R. Jackiw, Rev. Mod. Phys. **49**, 681 (1977).

[4] L. Landau et E. Lifshitz, *Mécanique Quantique*, Éditions MIR, Moscou 1966.

[5] M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions*, Dover Publications, INC, New York, (1964).

[6] F. Cooper, A. Khare and U. Sukhatme, Phys.Rep. **251**, 267, (1995).

[7] A. Khare and U. Sukhatme, J.Phys.A:Math. Gen.**21**,L501, (1988).

[8] K. Cahill, A. Comtet and R. Glauber, Phys.Lett. **B64**, 283 (1976).

[9] T. Weidig, hep-th/9912005.



Figure 1: The density potential $U(\phi)$ given by eq.(1).



Figure 2: The density potential $U(\phi)$ given by eq.(2).



Figure 3: The Schrodinger potential for model I



Figure 4: The Schrodinger potential for model II